

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin 17 (KRT17) (Basal Epithelial Marker) Antibody, IgG2b, Clone: [E3, same as Ks17.E3], Mouse, Monoclonal NBT-3872-MSM1-P1ABX

| | |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname | Cytokeratin 17 (KRT17) (Basal Epithelial Marker) Antibody, IgG2b, Clone: [E3, same as Ks17.E3], Mouse, Monoclonal |
| Artikelnummer | NBT-3872-MSM1-P1ABX |
| Hersteller Artikelnummer | 3872-MSM1-P1ABX |
| Alternativnummer | NBT-3872-MSM1-P1ABX-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC, WB |
| Spezies Reaktivität | Bovine, Goat, Human, Porcine, Rat |
| Immunogen | Cytoskeletal fraction of rat colon epithelium |
| Produktbeschreibung | Cytokeratin 17 (CK17) is normally expressed in the basal cells of complex epithelia but not in stratified or simple epithelia. Antibody to CK17 is an excellent tool to distinguish myoepithelial cells from luminal epithelium of various glandssuch as m... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [E3, same as Ks17.E3] |
| Molekulargewicht | 46kDa |
| Isotyp | IgG2b |

| | |
|------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| NCBI | 3872 |
| UniProt | Q04695 |
| Formulierung | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Western Blot (1-2ug/ml), Immunohistochemistry (Frozen & formalin-fixed) (1-2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with |