

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Cytokeratin 18 (KRT18) Antibody, IgG1, Clone: [C-04, same as Ks18.04], Mouse, Monoclonal NBT-3875-MSM7-P0**

Artikelname	Cytokeratin 18 (KRT18) Antibody, IgG1, Clone: [C-04, same as Ks18.04], Mouse, Monoclonal
Artikelnummer	NBT-3875-MSM7-P0
Hersteller Artikelnummer	3875-MSM7-P0
Alternativnummer	NBT-3875-MSM7-P0-20,NBT-3875-MSM7-P0-100
Hersteller	NeoBiotechnologies
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC, WB
Spezies Reaktivität	Bovine, Canine, Equine, Feline, Ferret, Goat, Hamster, Human, Monkey, Mouse, Porcine, Rat, Sheep
Immunogen	Cytoskeletal preparation extracted from the human vulvar squamous carcinoma cell line A431
Produktbeschreibung	This MAb reacts with a wide variety of simple epithelia. It does not react with stratified squamous epithelia. It reacts with epithelial tumors of the gastrointestinal tract, lung, breast, pancreas, ovary, and thyroid. Cytokeratin 18, which belongs t...
Klonalität	Monoclonal
Klon-Bezeichnung	[C-04, same as Ks18.04]
Molekulargewicht	45kDa

Isotyp	IgG1
NCBI	<a href="#">3875</a>
UniProt	<a href="#">P05783</a>
Formulierung	200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Antibody Type	Monoclonal Antibody
Anwendungsbeschreibung	Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Western Blot (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM ED