

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

MIF (Macrophage Migration Inhibitory Factor) Antibody, Clone: [MIF/6279], Mouse, Monoclonal NBT-4282-MSM9-P1ABX

| | |
|--------------------------|--|
| Artikelname | MIF (Macrophage Migration Inhibitory Factor) Antibody, Clone: [MIF/6279], Mouse, Monoclonal |
| Artikelnummer | NBT-4282-MSM9-P1ABX |
| Hersteller Artikelnummer | 4282-MSM9-P1ABX |
| Alternativnummer | NBT-4282-MSM9-P1ABX-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment of human MIF protein (exact sequence is proprietary) |
| Produktbeschreibung | Macrophage migration inhibitory factor, known as MIF or glycosylationinhibiting factor, is a secreted, homotrimeric, pro-inflammatory cytokine that modulates macrophage and T cell function and is an important regulator of host response to infection. ... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [MIF/6279] |
| Molekulargewicht | 13kDa |

| | |
|------------------------|--|
| NCBI | 4282 |
| UniProt | P14174 |
| Formulierung | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Western Blot (1-2ug/ml),,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 |