

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD45RB (B-Cell Marker) Antibody, IgG1, Clone: [BRA-11, same as BRA-11G], Mouse, Monoclonal NBT-5788-MSM6-P0**

|                          |   |
|--------------------------|---|
| Artikelname              | CD45RB (B-Cell Marker) Antibody, IgG1, Clone: [BRA-11, same as BRA-11G], Mouse, Monoclonal  |
| Artikelnummer            | NBT-5788-MSM6-P0  |
| Hersteller Artikelnummer | 5788-MSM6-P0  |
| Alternativnummer         | NBT-5788-MSM6-P0-20,NBT-5788-MSM6-P0-100  |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | FC, IF, IHC, WB   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Non-T, non-B CALLA positive ALL cell line REH (Leucocyte Workshop IV and V)   |
| Produktbeschreibung      | CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms ar... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [BRA-11, same as BRA-11G]   |
| Molekulargewicht         | 180-220kDa  |

|                        |   |
|------------------------|---|
| Isotyp                 | IgG1  |
| NCBI                   | <a href="#">5788</a>  |
| UniProt                | <a href="#">P08575</a>  |
| Formulierung           | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.  |
| Antibody Type          | Monoclonal Antibody   |
| Anwendungsbeschreibung | Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Western Blot (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM |