

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Actin, Smooth Muscle (Leiomyosarcoma Marker) Antibody, IgG2a, Clone: [1A4 + ACTA2/791], Mouse, Monoclonal NBT-59-MSM3-P1ABX**

|                          |   |
|--------------------------|---|
| Artikelname              | Actin, Smooth Muscle (Leiomyosarcoma Marker) Antibody, IgG2a, Clone: [1A4 + ACTA2/791], Mouse, Monoclonal   |
| Artikelnummer            | NBT-59-MSM3-P1ABX   |
| Hersteller Artikelnummer | 59-MSM3-P1ABX   |
| Alternativnummer         | NBT-59-MSM3-P1ABX-100   |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | FC, IF, IHC   |
| Spezies Reaktivität      | Human, Rat  |
| Immunogen                | N-Terminal decapeptide of alpha smooth muscle isoform of actin and conjugated to KLH (1A4), Recombinant full-length human ACTA2 protein (ACTA2/791)   |
| Produktbeschreibung      | Actin is a major component of the cytoskeleton and is present in most cell types. It is highly specific to actin from smooth muscles. This MAb does not stain cardiac or skeletal muscle, however, it does stain myofibroblasts and myoepithelial cells. T... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [1A4 + ACTA2/791]   |
| Molekulargewicht         | 42kDa   |

|                        |  |
|------------------------|--|
| Isotyp                 | IgG2a  |
| NCBI                   | <a href="#">59</a>   |
| UniProt                | <a href="#">P62736</a>   |
| Formulierung           | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.   |
| Antibody Type          | Monoclonal Antibody  |
| Anwendungsbeschreibung | Flow Cytometry (1-2ug/million cells), ,Immunofluorescence (1-2ug/ml), ,Immunohistochemistry (Formalin-fixed) (0.25-0.5ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for |