

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Recombinant Ribonucleotide Reductase M1 / RRM1 Antibody, Clone: [RRM1/4372R], Rabbit, Monoclonal NBT-6240-RBM1-P0**

|                          |   |
|--------------------------|---|
| Artikelname              | Recombinant Ribonucleotide Reductase M1 / RRM1 Antibody, Clone: [RRM1/4372R], Rabbit, Monoclonal  |
| Artikelnummer            | NBT-6240-RBM1-P0  |
| Hersteller Artikelnummer | 6240-RBM1-P0  |
| Alternativnummer         | NBT-6240-RBM1-P0-20,NBT-6240-RBM1-P0-100  |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Rabbit  |
| Kategorie                | Antikörper  |
| Applikation              | IHC, WB   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Synthetic peptide corresponding to residues within aa700-800 of human RRM1 (exact sequence is proprietary)  |
| Produktbeschreibung      | Ribonucleotide reductase M1 polypeptide (RRM1) is one of two non-identical subunits for ribonucleoside-diphosphate reductase, an enzyme which catalyzes the biosynthesis of deoxyribonucleotides from the corresponding ribonucleotides. It provides the p... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [RRM1/4372R]  |
| Molekulargewicht         | 94kDa   |

|                        |  |
|------------------------|--|
| NCBI                   | <a href="#">6240</a>   |
| UniProt                | <a href="#">P23921</a>   |
| Formulierung           | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.   |
| Antibody Type          | Recombinant Monoclonal Antibody  |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes),Optimal dilution fo |