

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

C1QA / Complement C1q A-Chain Antibody, IgG2b, Clone: [C1QA/2955], Mouse, Monoclonal NBT-712-MSM5-P0

| | |
|--------------------------|---|
| Artikelname | C1QA / Complement C1q A-Chain Antibody, IgG2b, Clone: [C1QA/2955], Mouse, Monoclonal |
| Artikelnummer | NBT-712-MSM5-P0 |
| Hersteller Artikelnummer | 712-MSM5-P0 |
| Alternativnummer | NBT-712-MSM5-P0-20,NBT-712-MSM5-P0-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment (around aa 104-237) of human C1QA protein (exact sequence is proprietary) |
| Produktbeschreibung | C1q, a subcomponent of the classical complement pathway, is composed of nine subunits that mediate classical complement activation and thereby play an important role in the immune response. Six of these subunits are disulfide-linked dimers of chains ... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [C1QA/2955] |
| Molekulargewicht | 29kDa |

| | |
|------------------------|---|
| Isotyp | IgG2b |
| NCBI | 712 |
| UniProt | P02745 |
| Formulierung | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Western Blot (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 min |