

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Ezrin / p81 Antibody, IgG2b, Clone: [CPTC-Ezrin-1], Mouse, Monoclonal NBT-7430-MSM1-P1ABX

| | |
|--------------------------|---|
| Artikelname | Ezrin / p81 Antibody, IgG2b, Clone: [CPTC-Ezrin-1], Mouse, Monoclonal |
| Artikelnummer | NBT-7430-MSM1-P1ABX |
| Hersteller Artikelnummer | 7430-MSM1-P1ABX |
| Alternativnummer | NBT-7430-MSM1-P1ABX-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human full-length Ezrin protein |
| Produktbeschreibung | Ezrin, Moesin and Radixin belong to a family of highly homologous actin-associated proteins that are localized just beneath the plasma membrane. The proteins are believed to be involved in the mediation of interactions between cytoskeletal and membra... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [CPTC-Ezrin-1] |
| Molekulargewicht | 87kDa |
| Isotyp | IgG2b |

| | |
|------------------------|---|
| NCBI | 7430 |
| UniProt | P15311 |
| Formulierung | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Western Blotting (1-2ug/ml), Immunohistology (Formalin-fixed) (1-2ug/ml for 30 min at RT),(Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate Buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes),Optima |