

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

TCL1 (T-Cell Marker) Antibody, IgG2a, Clone: [TCL1/2079], Mouse, Monoclonal NBT-8115-MSM2-P1ABX

| | |
|--------------------------|---|
| Artikelname | TCL1 (T-Cell Marker) Antibody, IgG2a, Clone: [TCL1/2079], Mouse, Monoclonal |
| Artikelnummer | NBT-8115-MSM2-P1ABX |
| Hersteller Artikelnummer | 8115-MSM2-P1ABX |
| Alternativnummer | NBT-8115-MSM2-P1ABX-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment (around aa 2-109) of human TCL1 protein (exact sequence is proprietary) |
| Produktbeschreibung | T-cell leukemia/Lymphoma Protein 1A (TCL-1A), also known as p14TCL1 is a product of the TCL1 gene that is involved in T-cell prolymphocytic leukemia (TPLL). T-PLL is a rare form of mature T-cell leukemia, which is consistently associated with chromos... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [TCL1/2079] |
| Molekulargewicht | 14kDa |

| | |
|------------------------|---|
| Isotyp | IgG2a |
| NCBI | 10232 |
| UniProt | P56279 |
| Formulierung | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Western Blot (1-2ug/ml),Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minu |