

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Sarcomeric Actinin Alpha 2 / ACTN2 Antibody, IgG2b, Clone: [ACTN2/3292], Mouse, Monoclonal NBT-88-MSM2-P1ABX**

|                          |   |
|--------------------------|---|
| Artikelname              | Sarcomeric Actinin Alpha 2 / ACTN2 Antibody, IgG2b, Clone: [ACTN2/3292], Mouse, Monoclonal  |
| Artikelnummer            | NBT-88-MSM2-P1ABX   |
| Hersteller Artikelnummer | 88-MSM2-P1ABX   |
| Alternativnummer         | NBT-88-MSM2-P1ABX-100   |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC, WB   |
| Spezies Reaktivität      | Human   |
| Immunogen                | A recombinant fragment (aa557-692) of human ACTN2 protein (exact sequence is proprietary)   |
| Produktbeschreibung      | Alpha actinins belong to the spectrin gene superfamily which represents a diverse group of cytoskeletal proteins, including the alpha and beta spectrins and dystrophins. Alpha actinin is an actin-binding protein with multiple roles in different cell ... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [ACTN2/3292]  |
| Molekulargewicht         | 103kDa  |

|                        |  |
|------------------------|--|
| Isotyp                 | IgG2b  |
| NCBI                   | <a href="#">88</a>   |
| UniProt                | <a href="#">P35609</a>   |
| Formulierung           | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.   |
| Antibody Type          | Monoclonal Antibody  |
| Anwendungsbeschreibung | Western Blotting (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT), (Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes) |