

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Recombinant CD3e (T-Cell Marker) Antibody, IgG1, Clone: [rC3e/1931], Mouse, Monoclonal NBT-916-MSM20-P1ABX**

|                          |   |
|--------------------------|---|
| Artikelname              | Recombinant CD3e (T-Cell Marker) Antibody, IgG1, Clone: [rC3e/1931], Mouse, Monoclonal  |
| Artikelnummer            | NBT-916-MSM20-P1ABX   |
| Hersteller Artikelnummer | 916-MSM20-P1ABX   |
| Alternativnummer         | NBT-916-MSM20-P1ABX-100   |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | ELISA, IHC, WB  |
| Spezies Reaktivität      | Human   |
| Immunogen                | A recombinant human CD3e protein fragment (around aa 23-119) (exact sequence is proprietary)  |
| Produktbeschreibung      | Recognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [rC3e/1931]   |
| Molekulargewicht         | 20kDa   |

|                        |   |
|------------------------|---|
| Isotyp                 | IgG1  |
| NCBI                   | <a href="#">916</a>   |
| UniProt                | <a href="#">P07766</a>  |
| Formulierung           | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.  |
| Antibody Type          | Recombinant Monoclonal Antibody   |
| Anwendungsbeschreibung | ELISA (Use Ab at 2-4ug/ml for coating) (Order Ab without BSA), Western Blot (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA |