

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD7 (T-Cell Leukemia Marker) Antibody, IgG1, Clone: [CD7/3737], Mouse, Monoclonal NBT-924-MSM7-P0

| | |
|--------------------------|---|
| Artikelname | CD7 (T-Cell Leukemia Marker) Antibody, IgG1, Clone: [CD7/3737], Mouse, Monoclonal |
| Artikelnummer | NBT-924-MSM7-P0 |
| Hersteller Artikelnummer | 924-MSM7-P0 |
| Alternativnummer | NBT-924-MSM7-P0-20,NBT-924-MSM7-P0-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment (around aa 72-175) of human CD7 protein (exact sequence is proprietary) |
| Produktbeschreibung | Recognizes a protein of 40kDa, identified as CD7 (also known as gp40, Leu9). CD7 is a member of the immunoglobulin gene superfamily. Its N-terminal amino acids 1-107 are highly homologous to Ig kappa-L chains whereas the carboxyl-terminal region of t... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [CD7/3737] |
| Molekulargewicht | 40kDa |

| | |
|------------------------|--|
| Isotyp | IgG1 |
| NCBI | 924 |
| UniProt | P09564 |
| Formulierung | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Western Blot (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT), (Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 2 |