

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD9 (TSPAN29) (Motility-Related Protein-1) Antibody, IgG1, Clone: [CD9/2343], Mouse, Monoclonal NBT-928-MSM3-P0

| | |
|--------------------------|---|
| Artikelname | CD9 (TSPAN29) (Motility-Related Protein-1) Antibody, IgG1, Clone: [CD9/2343], Mouse, Monoclonal |
| Artikelnummer | NBT-928-MSM3-P0 |
| Hersteller Artikelnummer | 928-MSM3-P0 |
| Alternativnummer | NBT-928-MSM3-P0-20,NBT-928-MSM3-P0-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human CD9 protein |
| Produktbeschreibung | CD9 is a type IV transmembrane glycoprotein with four transmembrane domains. CD9 on pre-B cells may play a role in cell-cell adhesion. In addition, CD9 may play a role in signal transduction mediated by interaction with low molecular weight GTP bindi... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [CD9/2343] |
| Molekulargewicht | 24kDa |
| Isotyp | IgG1 |

| | |
|------------------------|---|
| NCBI | 928 |
| UniProt | P21926 |
| Formulierung | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes),Optimal dilution f |