

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant CD68 (Macrophage Marker) Antibody, Clone: [C68/2908R], Rabbit, Monoclonal NBT-968-RBM10-P0

| | |
|--------------------------|---|
| Artikelname | Recombinant CD68 (Macrophage Marker) Antibody, Clone: [C68/2908R], Rabbit, Monoclonal |
| Artikelnummer | NBT-968-RBM10-P0 |
| Hersteller Artikelnummer | 968-RBM10-P0 |
| Alternativnummer | NBT-968-RBM10-P0-20,NBT-968-RBM10-P0-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human LAMP4 / CD68 protein |
| Produktbeschreibung | This antibody recognizes a glycoprotein of 110kDa, which is identified as CD68. It is important for identifying macrophages in tissue sections. It stains macrophages in a wide variety of human tissues, including Kupffer cells and macrophages in the r... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [C68/2908R] |
| Molekulargewicht | 110kDa |
| NCBI | 968 |

| | |
|------------------------|--|
| UniProt | P34810 |
| Formulierung | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Anwendungsbeschreibung | Western Blot (1-2ug/ml),,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 2 |