

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### HDAC1 (C-terminus) Antibody, Rabbit, Polyclonal ROC-600-401-879S

|                          |                                                                                                                                                                                                      |
|--------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | HDAC1 (C-terminus) Antibody, Rabbit, Polyclonal                                                                                                                                                      |
| Artikelnummer            | ROC-600-401-879S                                                                                                                                                                                     |
| Hersteller Artikelnummer | 600-401-879S                                                                                                                                                                                         |
| Alternativnummer         | ROC-600-401-879S                                                                                                                                                                                     |
| Hersteller               | Rockland Immunochemicals                                                                                                                                                                             |
| Wirt                     | Rabbit                                                                                                                                                                                               |
| Kategorie                | Antikörper                                                                                                                                                                                           |
| Applikation              | ELISA, IF, IHC, Multiplex Assay, WB                                                                                                                                                                  |
| Spezies Reaktivität      | Human                                                                                                                                                                                                |
| Immunogen                | Anti-HDAC-1 antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to a C-Terminal region near amino acids 450-482 of Human HDAC-1. |
| Konjugation              | Unconjugated                                                                                                                                                                                         |
| Klonalität               | Polyclonal                                                                                                                                                                                           |
| Konzentration            | 1.0 mg/mL by UV absorbance at 280 nm                                                                                                                                                                 |
| NCBI                     | <a href="#">3065</a>                                                                                                                                                                                 |
| UniProt                  | <a href="#">Q13547</a>                                                                                                                                                                               |
| Puffer                   | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2                                                                                                                                           |

|                        |                                                                                                                                                                                                                                                                |
|------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Formulierung           | Liquid (sterile filtered)                                                                                                                                                                                                                                      |
| Target-Kategorie       | Human                                                                                                                                                                                                                                                          |
| Antibody Type          | Primary Antibody                                                                                                                                                                                                                                               |
| Application Verdünnung | ELISA: 1:10,000 - 1:50,000, IHC: 1:200 - 1:1,000, IF Microscopy: 10ug / ml, WB: 1:1,000 - 1:5,000                                                                                                                                                              |
| Anwendungsbeschreibung | Anti-HDAC-1 Antibody has been tested for use in ELISA, immunohistochemistry, immunofluorescence, and western blot. Specific conditions for reactivity should be optimized by the end user. Specific nuclear staining is observed by IHC. Expect bands at 65 kD |