

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Goat F(ab)2 anti-Mouse IgG (Fc)-RPE, MinX Bo,Ho,Hu, Polyclonal ROC-710-1822

|                                  |                                                                                                        |
|----------------------------------|--------------------------------------------------------------------------------------------------------|
| Artikelname                      | Goat F(ab)2 anti-Mouse IgG (Fc)-RPE, MinX Bo,Ho,Hu, Polyclonal                                         |
| Artikelnummer                    | ROC-710-1822                                                                                           |
| Hersteller Artikelnummer         | 710-1822                                                                                               |
| Alternativnummer                 | ROC-710-1822                                                                                           |
| Hersteller                       | Rockland Immunochemicals                                                                               |
| Wirt                             | Goat                                                                                                   |
| Kategorie                        | Antikörper                                                                                             |
| Applikation                      | FACS,IF                                                                                                |
| Spezies Reaktivität              | Mouse                                                                                                  |
| Immunogen                        | F(ab)2 anti-Mouse IgG F(c) was produced by repeated immunization with Mouse IgG F(c) fragment in goat. |
| Konjugation                      | RPE                                                                                                    |
| Format                           | F(ab')2                                                                                                |
| Spezifität                       | IgG (Fc)                                                                                               |
| Minimale Kreuzreaktivität (MinX) | Bovine,Equine,Human                                                                                    |
| Klonalität                       | Polyclonal                                                                                             |
| Konzentration                    | 0.5 mg/mL by absorbance = 82.0 at 565 nm                                                               |
| Puffer                           | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2                                             |

|                        |                                                                                                                                                                                                                                                                |
|------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Formulierung           | Lyophilized                                                                                                                                                                                                                                                    |
| Target-Kategorie       | Mouse                                                                                                                                                                                                                                                          |
| Antibody Type          | Secondary Antibody                                                                                                                                                                                                                                             |
| Application Verdünnung | Flow Cytometry: 1:100 - 1:250, IF Microscopy: 1:100 - 1:250                                                                                                                                                                                                    |
| Anwendungsbeschreibung | Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maxi |