

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

flt3-Ligand Antibody, Unconjugated, Rabbit, Polyclonal PRS-XP-5138

Artikelname	flt3-Ligand Antibody, Unconjugated, Rabbit, Polyclonal
Artikelnummer	PRS-XP-5138
Hersteller Artikelnummer	XP-5138
Alternativnummer	PRS-XP-5138-0.1
Hersteller	ProSci
Wirt	Rabbit
Kategorie	Antikörper
Applikation	ELISA, NeA, WB
Spezies Reaktivität	Human
Immunogen	Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hflt3-Ligand.
Konjugation	Unconjugated
Klonalität	Polyclonal
Konzentration	batch dependent
NCBI	2323
UniProt	P49771
Formulierung	Lyophilized
Application Verdünnung	Centrifuge vial prior to opening.

Anwendungsbeschreibung	Neutralization: To yield one-half maximal inhibition [ND50] of the biological activity of hflt3- Ligand (15 ng/mL), a concentration of 0.25 - 0.75 µg/mL of this antibody is required. ELISA: To detect hflt3- Ligand by direct ELISA (using 100 µL/well antibody solution) a concentration of at least 0.5 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of 0.2 - 0.4 ng/well of recombinant hflt 3-Ligand. Sandwich: To detect hFGF-basic by sandwich ELISA (using 100 µL/well antibody solution) a concentration of 0.25 - 1.0 µg/mL of this antibody is required. This biotinylated polyclonal antibody, in conjunction with our Polyclonal Anti-Human FGF-basic (XP-5138Bt) as a capture antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hFGF-basic. Western Blot: To detect hflt3-Ligand by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant hflt3-Ligand is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.
------------------------	--