

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

**CD56 / NCAM(123A8, same as 56C04), CF405S conjugate, 0.1mg/mL,  
Clone: [123A8 same as 56C04], Mouse, Monoclonal  
BOT-BNC040692-500**

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | CD56 / NCAM(123A8, same as 56C04), CF405S conjugate, 0.1mg/mL,<br>Clone: [123A8 same as 56C04], Mouse, Monoclonal                                                                                                                                             |
| Artikelnummer            | BOT-BNC040692-500                                                                                                                                                                                                                                             |
| Hersteller Artikelnummer | BNC040692-500                                                                                                                                                                                                                                                 |
| Alternativnummer         | BOT-BNC040692-500-500UL                                                                                                                                                                                                                                       |
| Hersteller               | Biotium                                                                                                                                                                                                                                                       |
| Wirt                     | Mouse                                                                                                                                                                                                                                                         |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | IHC                                                                                                                                                                                                                                                           |
| Spezies Reaktivität      | Human                                                                                                                                                                                                                                                         |
| Immunogen                | Membrane preparation of a small cell lung carcinoma                                                                                                                                                                                                           |
| Konjugation              | CF405S                                                                                                                                                                                                                                                        |
| Produktbeschreibung      | This MAb reacts with an extracellular domain (close to transmembrane) of CD56/NCAM. Three isoforms of neural cell adhesion molecule (NCAM) are produced by differential splicing of the RNA transcript from a single gene. The 135 kDa isoform is the basi... |
| Klonalität               | Monoclonal                                                                                                                                                                                                                                                    |
| Konzentration            | 0.1 mg/mL                                                                                                                                                                                                                                                     |

|                        |                                                                                                                                                                                                                                                                                                            |
|------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Klon-Bezeichnung       | [123A8 same as 56C04]                                                                                                                                                                                                                                                                                      |
| Molekulargewicht       | 180, 145 and 125 kDa                                                                                                                                                                                                                                                                                       |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide                                                                                                                                                                                                                                                                                 |
| Quelle                 | Animal                                                                                                                                                                                                                                                                                                     |
| Anwendungsbeschreibung | Immunohistology formalin-fixed 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Immunofluorescence 1-2 ug/mL Optimal dilution for a specific application should be determined by user |