

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

IgG (Immunoglobulin Gamma Heavy Chain) (B-Cell Marker), CF405S conjugate, 0.1mg/mL, Rabbit BOT-BNC040728-500

Artikelname	IgG (Immunoglobulin Gamma Heavy Chain) (B-Cell Marker), CF405S conjugate, 0.1mg/mL, Rabbit
Artikelnummer	BOT-BNC040728-500
Hersteller Artikelnummer	BNC040728-500
Alternativnummer	BOT-BNC040728-500-500UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Purified human Ig Gamma Chain
Konjugation	CF405S
Produktbeschreibung	This antibody recognizes a protein of 75 kDa, identified as gamma heavy chain of human immunoglobulins. It does not cross-react with alpha (IgA), mu (IgM), epsilon (IgE), or delta (IgD), heavy chains, T-cells, monocytes, granulocytes, or erythrocytes...
Konzentration	0.1 mg/mL
Molekulargewicht	75 kDa
UniProt	P01861 , P01860 , P01859 , P01857

Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user