

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

N-Cadherin / Cadherin-2 / CD325 (NCAD) (8C11), CF405S conjugate, 0.1mg/mL, Clone: [8C11], Mouse, Monoclonal BOT-BNC041426-100

Artikelname	N-Cadherin / Cadherin-2 / CD325 (NCAD) (8C11), CF405S conjugate, 0.1mg/mL, Clone: [8C11], Mouse, Monoclonal
Artikelnummer	BOT-BNC041426-100
Hersteller Artikelnummer	BNC041426-100
Alternativnummer	BOT-BNC041426-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human, Mouse
Immunogen	Recombinant human N-cadherin extracellular domain (exact sequence is proprietary)
Konjugation	CF405S
Produktbeschreibung	This antibody recognizes a protein of ~140 kDa, identified as N-Cadherin (NCAD), also known as CD325. NCAD is a member of the Cadherin superfamily, and consists of five extracellular repeats, a transmembrane domain and a cytoplasmic domain. CD325 def...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[8C11]
Molekulargewicht	130-140 kDa
UniProt	P19022
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user