

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD117 / c-Kit (Marker for Gastrointestinal Stromal Tumors)(KIT/2674), CF405S conjugate, 0.1mg/mL, Clone: [KIT/2674], Mouse, Monoclonal BOT-BNC042674-500

| | |
|--------------------------|---|
| Artikelname | CD117 / c-Kit (Marker for Gastrointestinal Stromal Tumors)(KIT/2674), CF405S conjugate, 0.1mg/mL, Clone: [KIT/2674], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC042674-500 |
| Hersteller Artikelnummer | BNC042674-500 |
| Alternativnummer | BOT-BNC042674-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human KIT protein |
| Konjugation | CF405S |
| Produktbeschreibung | CD117, also known as c-kit, is a member of the immunoglobulin superfamily. It is a receptor tyrosine kinase that is expressed on hematopoietic progenitor cells. It binds to the c-kit ligand KITLG/SCF and has roles in cell survival and cell proliferatio... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [KIT/2674] |

| | |
|------------------------|--|
| Molekulargewicht | 145 kDa |
| UniProt | P10721 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |