

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Fascin-1(FSCN1/416), CF640R conjugate, 0.1mg/mL, Clone: [FSCN1/416], Mouse, Monoclonal BOT-BNC400416-500

| | |
|--------------------------|---|
| Artikelname | Fascin-1(FSCN1/416), CF640R conjugate, 0.1mg/mL, Clone: [FSCN1/416], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC400416-500 |
| Hersteller Artikelnummer | BNC400416-500 |
| Alternativnummer | BOT-BNC400416-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Full length recombinant human FSCN1 protein |
| Konjugation | CF640R |
| Produktbeschreibung | Recognizes a protein of 55 kDa, which is identified as fascin-1. Its actin binding ability is regulated by phosphorylation. Antibody to fascin-1 is a very sensitive marker for Reed-Sternberg cells and variants in nodular sclerosis, mixed cellularity,... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [FSCN1/416] |

| | |
|------------------------|---|
| Molekulargewicht | 55 kDa |
| UniProt | Q16658 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin) Staining of formalin-fixed tissues requires boiling tissue sections in 1 mM EDTA, pH 8.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |