

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

ODC-1 (Ornithine Decarboxylase-1)(ODC1/485), CF640R conjugate, 0.1mg/mL, Clone: [ODC1/485], Mouse, Monoclonal BOT-BNC400485-100

| | |
|--------------------------|---|
| Artikelname | ODC-1 (Ornithine Decarboxylase-1)(ODC1/485), CF640R conjugate, 0.1mg/mL, Clone: [ODC1/485], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC400485-100 |
| Hersteller Artikelnummer | BNC400485-100 |
| Alternativnummer | BOT-BNC400485-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC, WB |
| Spezies Reaktivität | Human, Rat |
| Immunogen | Recombinant human ODC-1 protein |
| Konjugation | CF640R |
| Produktbeschreibung | Recognizes a 53 kDa protein, identified as the Ornithine Decarboxylase (ODC-1). ODC is the initial and rate-limiting enzyme in the biosynthetic pathway of polyamines and is involved in the conversion of ornithine to putrescine. The biological activit... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [ODC1/485] |

| | |
|------------------------|--|
| Molekulargewicht | 53 kDa |
| UniProt | P11926 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |