

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **von Willebrand Factor(3E2D10 + VWF635), CF640R conjugate, 0.1mg/mL, Clone: [3E2D10 VWF635], Mouse, Monoclonal BOT-BNC400646-500**

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | von Willebrand Factor(3E2D10 + VWF635), CF640R conjugate, 0.1mg/mL, Clone: [3E2D10 VWF635], Mouse, Monoclonal                                                                                                                                                 |
| Artikelnummer            | BOT-BNC400646-500                                                                                                                                                                                                                                             |
| Hersteller Artikelnummer | BNC400646-500                                                                                                                                                                                                                                                 |
| Alternativnummer         | BOT-BNC400646-500-500UL                                                                                                                                                                                                                                       |
| Hersteller               | Biotium                                                                                                                                                                                                                                                       |
| Wirt                     | Mouse                                                                                                                                                                                                                                                         |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | IHC                                                                                                                                                                                                                                                           |
| Spezies Reaktivität      | Human                                                                                                                                                                                                                                                         |
| Immunogen                | Recombinant human vWF fragment                                                                                                                                                                                                                                |
| Konjugation              | CF640R                                                                                                                                                                                                                                                        |
| Produktbeschreibung      | von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslation... |
| Klonalität               | Monoclonal                                                                                                                                                                                                                                                    |
| Konzentration            | 0.1 mg/mL                                                                                                                                                                                                                                                     |
| Klon-Bezeichnung         | [3E2D10 VWF635]                                                                                                                                                                                                                                               |

|                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
|------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Molekulargewicht       | 250 kDa                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| UniProt                | <a href="#">P04275</a>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
| Quelle                 | Animal                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunoprecipitation: 0.5-1 ug/500 ug protein lysate Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1.0 ug/mL Optimal dilution for a specific application should be determined by user</p> |