

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

DNMT3A / DNA Methyltransferase 3 Alpha (PCRP-DNMT3A-1E2), CF640R conjugate, 0.1mg/mL, IgG1, Clone: [PCRP-DNMT3A-1E2], Mouse, Monoclonal BOT-BNC401898-500

| | |
|--------------------------|---|
| Artikelname | DNMT3A / DNA Methyltransferase 3 Alpha (PCRP-DNMT3A-1E2), CF640R conjugate, 0.1mg/mL, IgG1, Clone: [PCRP-DNMT3A-1E2], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC401898-500 |
| Hersteller Artikelnummer | BNC401898-500 |
| Alternativnummer | BOT-BNC401898-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human DNMT3A protein |
| Konjugation | CF640R |
| Produktbeschreibung | The specificity of this monoclonal antibody to its intended target was validated by HuProt(TM)Array, containing more than 19,000, full-length human proteins. DNMT3A is required for genome-wide de novo methylation and is essential for the establishmen... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|---|
| Klon-Bezeichnung | [PCRP-DNMT3A-1E2] |
| Molekulargewicht | 102 kDa |
| Isotyp | IgG1 |
| UniProt | Q16655 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Immunoprecipitation 2-4 ug/mg protein Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Western blotting 0.5-1.0 ug/mL Optimal dilution for a specific application should be determined by user |