

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

PAX5 / BSAP (Early B-Cell Marker) (PCRP-PAX5-1B7), CF640R conjugate, 0.1mg/mL, IgG1, Clone: [PCRP-PAX5-1B7], Mouse, Monoclonal BOT-BNC401908-100

| | |
|--------------------------|---|
| Artikelname | PAX5 / BSAP (Early B-Cell Marker) (PCRP-PAX5-1B7), CF640R conjugate, 0.1mg/mL, IgG1, Clone: [PCRP-PAX5-1B7], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC401908-100 |
| Hersteller Artikelnummer | BNC401908-100 |
| Alternativnummer | BOT-BNC401908-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human PAX5 protein |
| Konjugation | CF640R |
| Produktbeschreibung | The specificity of this monoclonal antibody to its intended target was validated by HuProt(TM) Array, containing more than 19,000, full-length human proteins. The PAX5 gene is a member of the paired box (PAX) family of transcription factors. The cent... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [PCRP-PAX5-1B7] |

| | |
|------------------------|--|
| Molekulargewicht | 42 kDa |
| Isotyp | IgG1 |
| UniProt | Q02548 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |