

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **TIMP1 (Colorectal Cancer Biomarker / Marker of Lymph Node Metastasis)(rTIMP1/1710), CF640R conjugate, 0.1mg/mL, Clone: [rTIMP1/1710], Mouse, Monoclonal BOT-BNC401943-500**

Artikelname	TIMP1 (Colorectal Cancer Biomarker / Marker of Lymph Node Metastasis)(rTIMP1/1710), CF640R conjugate, 0.1mg/mL, Clone: [rTIMP1/1710], Mouse, Monoclonal
Artikelnummer	BOT-BNC401943-500
Hersteller Artikelnummer	BNC401943-500
Alternativnummer	BOT-BNC401943-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human TIMP1 protein
Konjugation	CF640R
Produktbeschreibung	TIMP-1, TIMP-2, TIMP-3 and TIMP-4 (for tissue inhibitor of metalloproteinases -1, -2, -3 and -4) complex with metalloproteinases such as collagenases, gelatinases and stromelysins, resulting in irreversible inactivation of the metalloproteinase. TIMP...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[rTIMP1/1710]
Molekulargewicht	23 kDa
UniProt	<a href="#">P01033</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry