

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Cytokeratin 8 (KRT8) (rB22.1), CF640R conjugate, 0.1mg/mL, IgG1, Clone: [rB22.1], Mouse, Monoclonal BOT-BNC402175-500**

Artikelname	Cytokeratin 8 (KRT8) (rB22.1), CF640R conjugate, 0.1mg/mL, IgG1, Clone: [rB22.1], Mouse, Monoclonal
Artikelnummer	BOT-BNC402175-500
Hersteller Artikelnummer	BNC402175-500
Alternativnummer	BOT-BNC402175-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Bovine, Human, Porcine, Rabbit, Sheep
Immunogen	Recombinant full-length human KRT8 protein
Konjugation	CF640R
Produktbeschreibung	Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). CK8 is primarily found in the non-squamous epithelia and is present in majority of adenoc...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[rB22.1]

Molekulargewicht	52.5 kDa
Isotyp	IgG1
UniProt	<a href="#">P05787</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Does not react with mouse, rat, chicken, or <i>Xenopus laevis</i>  Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user