

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CFTR (Cystic Fibrosis Transmembrane Conductance Regulator) (CFTR/2290R), CF640R conjugate, 0.1mg/mL, Clone: [CFTR/2290R], Rabbit, Monoclonal BOT-BNC402290-500

| | |
|--------------------------|---|
| Artikelname | CFTR (Cystic Fibrosis Transmembrane Conductance Regulator) (CFTR/2290R), CF640R conjugate, 0.1mg/mL, Clone: [CFTR/2290R], Rabbit, Monoclonal |
| Artikelnummer | BOT-BNC402290-500 |
| Hersteller Artikelnummer | BNC402290-500 |
| Alternativnummer | BOT-BNC402290-500-500UL |
| Hersteller | Biotium |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human, Mouse |
| Immunogen | Recombinant full-length human CFTR protein |
| Konjugation | CF640R |
| Produktbeschreibung | This antibody recognizes a protein of 165-170 kDa, identified as cystic fibrosis transmembrane conductance regulator (CFTR). CFTR is composed of two membrane-spanning domains (MSD), two nucleotide-binding domains (NBD), and an R domain. It is structu... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [CFTR/2290R] |
| Molekulargewicht | 165-170 kDa |
| UniProt | P13569 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1.0 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |