

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Smooth Muscle Myosin Heavy Chain (SM-MHC) (MYH11/2303R), CF640R conjugate, 0.1mg/mL, Clone: [MYH11/2303R], Rabbit, Monoclonal BOT-BNC402303-500

|                          |   |
|--------------------------|---|
| Artikelname              | Smooth Muscle Myosin Heavy Chain (SM-MHC) (MYH11/2303R), CF640R conjugate, 0.1mg/mL, Clone: [MYH11/2303R], Rabbit, Monoclonal   |
| Artikelnummer            | BOT-BNC402303-500   |
| Hersteller Artikelnummer | BNC402303-500   |
| Alternativnummer         | BOT-BNC402303-500-500UL   |
| Hersteller               | Biotium   |
| Wirt                     | Rabbit  |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Bovine, Canine, Feline, Gallus, Guinea pig, Human, Porcine, Rabbit, Rat   |
| Immunogen                | Recombinant full-length human MYH11 protein   |
| Konjugation              | CF640R  |
| Produktbeschreibung      | Smooth muscle myosin heavy chain (SM-MHC) is a cytoplasmic structural protein, which is a major component of the contractile apparatus in smooth muscle cells. Expression of smooth muscle myosin is developmentally regulated, appearing early in smooth ... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |

|                        |   |
|------------------------|---|
| Klon-Bezeichnung       | [MYH11/2303R]   |
| Molekulargewicht       | 205 kDa (MHC-1) and 200 kDa (MHC-2)   |
| UniProt                | <a href="#">P35749</a>  |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide  |
| Quelle                 | Animal  |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |