

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Calcineurin B / PPP3R1(BE2.1), CF640R conjugate, 0.1mg/mL, Clone: [BE2.1], Mouse, Monoclonal BOT-BNC402321-100

| | |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname | Calcineurin B / PPP3R1(BE2.1), CF640R conjugate, 0.1mg/mL, Clone: [BE2.1], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC402321-100 |
| Hersteller Artikelnummer | BNC402321-100 |
| Alternativnummer | BOT-BNC402321-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Spezies Reaktivität | Human, Xenopus |
| Immunogen | Recombinant full-length human Calcineurin B protein |
| Konjugation | CF640R |
| Produktbeschreibung | This antibody recognizes the Calcineurin B alpha protein, encoded by the PPP3R1 (Protein Phosphatase 3 Regulatory Subunit B Alpha) gene. Calcineurin is an enzyme that dephosphorylates serine and threonine residues in proteins. It is a heterodimer of ... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [BE2.1] |
| Molekulargewicht | 19 kDa |

| | |
|------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| UniProt | P63098 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |