

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD25/ IL2RA (Activated Lymphocyte Marker) (IL2RA/2395), CF640R conjugate, 0.1mg/mL, IgG2b, Clone: [IL2RA/2395], Mouse, Monoclonal BOT-BNC402395-500

Artikelname	CD25/ IL2RA (Activated Lymphocyte Marker) (IL2RA/2395), CF640R conjugate, 0.1mg/mL, IgG2b, Clone: [IL2RA/2395], Mouse, Monoclonal
Artikelnummer	BOT-BNC402395-500
Hersteller Artikelnummer	BNC402395-500
Alternativnummer	BOT-BNC402395-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinantfragment of human IL2RA protein (around aa 42-183) (exact sequence is proprietary)
Konjugation	CF640R
Produktbeschreibung	This antibody recognizes a protein of 55 kDa, identified as CD25. It is expressed on activated T- and B-cells and activated monocytes/macrophages. With respect to lymphomas, CD25 is present on malignant cells of Hodgkins disease, HTLV-1 adult T-cell ...
Klonalität	Monoclonal

Konzentration	0.1 mg/mL
Klon-Bezeichnung	[IL2RA/2395]
Molekulargewicht	55 kDa
Isotyp	IgG2b
UniProt	P01589
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 2-4 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user