

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

p63 (Squamous, Basal & Myoepithelial Cell Marker) (TP63/2428), CF647 conjugate, 0.1mg/mL, Clone: [TP63/2428], Mouse, Monoclonal BOT-BNC472428-500

| | |
|--------------------------|---|
| Artikelname | p63 (Squamous, Basal & Myoepithelial Cell Marker) (TP63/2428), CF647 conjugate, 0.1mg/mL, Clone: [TP63/2428], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC472428-500 |
| Hersteller Artikelnummer | BNC472428-500 |
| Alternativnummer | BOT-BNC472428-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human p63 protein |
| Konjugation | CF647 |
| Produktbeschreibung | p63 is a homolog of the tumor suppressor p53. It is identified in basal cells in the epithelial layers of a variety of tissues, including epidermis, cervix, urothelium, breast and prostate. p63 was detected in nuclei of the basal epithelium in normal... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [TP63/2428] |

| | |
|------------------------|---|
| Molekulargewicht | 63 kDa |
| UniProt | Q9H3D4 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |