

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Cig2(3A11/5), CF647 conjugate, 0.1mg/mL, Clone: [3A11/5], Mouse, Monoclonal BOT-BNC472827-100**

Artikelname	Cig2(3A11/5), CF647 conjugate, 0.1mg/mL, Clone: [3A11/5], Mouse, Monoclonal
Artikelnummer	BOT-BNC472827-100
Hersteller Artikelnummer	BNC472827-100
Alternativnummer	BOT-BNC472827-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Spezies Reaktivität	Yeast
Immunogen	Bacterial <i>S. pombe</i> Cig2 Purified protein.
Konjugation	CF647
Produktbeschreibung	Cig2 is a cyclin that functions as the catalytic subunit of cyclin-dependent kinases (Cdks) in the fission yeast <i>Schizosaccharomyces pombe</i> . It is a B-type, S phase cyclin that is required for both the G1/S and G2/M cell cycle transitions and is expre...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[3A11/5]
Molekulargewicht	~50 kDa

UniProt	Not Known
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry