

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Golgi Complex (Marker for Human Cells) (GLG1/2829R), CF647 conjugate, 0.1mg/mL, Clone: [GLG1/2829R], Rabbit, Monoclonal BOT-BNC472829-100

Artikelname	Golgi Complex (Marker for Human Cells) (GLG1/2829R), CF647 conjugate, 0.1mg/mL, Clone: [GLG1/2829R], Rabbit, Monoclonal
Artikelnummer	BOT-BNC472829-100
Hersteller Artikelnummer	BNC472829-100
Alternativnummer	BOT-BNC472829-100-100UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	SU-DHL-1 large cell lymphoma cells.
Konjugation	CF647
Produktbeschreibung	This MAb recognizes an antigen associated with the Golgi complex in human cells only. It can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffuse staining ...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[GLG1/2829R]

Molekulargewicht	Not Known
UniProt	Not Known
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Does not react with mouse or rat Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user