

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Secretory Component / ECM1 (ECM1/2889R), CF647 conjugate, 0.1mg/mL, Clone: [ECM1/2889R], Rabbit, Monoclonal BOT-BNC472889-100

| | |
|--------------------------|---|
| Artikelname | Secretory Component / ECM1 (ECM1/2889R), CF647 conjugate, 0.1mg/mL, Clone: [ECM1/2889R], Rabbit, Monoclonal |
| Artikelnummer | BOT-BNC472889-100 |
| Hersteller Artikelnummer | BNC472889-100 |
| Alternativnummer | BOT-BNC472889-100-100UL |
| Hersteller | Biotium |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human, Rat |
| Immunogen | Recombinant full-length human ECM1 protein |
| Konjugation | CF647 |
| Produktbeschreibung | This MAb reacts with a reduction-resistant epitope present in both free and SIgA bound Secretory Component. It does not react with the cell lines lacking secretory component. The antibody is useful for studying the distribution and level of both free... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [ECM1/2889R] |

| | |
|------------------------|---|
| Molekulargewicht | ~80 kDa |
| UniProt | Q16610 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |