

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD68 (Macrophage Marker) (C68/2908R), CF647 conjugate, 0.1mg/mL, Clone: [C68/2908R], Rabbit, Monoclonal BOT-BNC472908-100**

Artikelname	CD68 (Macrophage Marker) (C68/2908R), CF647 conjugate, 0.1mg/mL, Clone: [C68/2908R], Rabbit, Monoclonal
Artikelnummer	BOT-BNC472908-100
Hersteller Artikelnummer	BNC472908-100
Alternativnummer	BOT-BNC472908-100-100UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human LAMP4 protein
Konjugation	CF647
Produktbeschreibung	This antibody recognizes a glycoprotein of 110 kDa, which is identified as CD68. It is important for identifying macrophages in tissue sections. It stains macrophages in a wide variety of human tissues, including Kupffer cells and macrophages in the ...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[C68/2908R]

Molekulargewicht	~110 kDa
UniProt	<a href="#">P34810</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer pH 6.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user