

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **HLA-Pan (MHC II) (HLA-Pan/2967R), CF647 conjugate, 0.1mg/mL, Clone: [HLA-Pan/2967R], Rabbit, Monoclonal BOT-BNC472967-100**

Artikelname	HLA-Pan (MHC II) (HLA-Pan/2967R), CF647 conjugate, 0.1mg/mL, Clone: [HLA-Pan/2967R], Rabbit, Monoclonal
Artikelnummer	BOT-BNC472967-100
Hersteller Artikelnummer	BNC472967-100
Alternativnummer	BOT-BNC472967-100-100UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	FC, IF, IHC, WB
Spezies Reaktivität	Human
Immunogen	Non-T, non-B human acute lymphoblastic leukemia REH6 cell line
Konjugation	CF647
Produktbeschreibung	This antibody reacts with a common epitope of human major histocompatibility (MHC) class II antigens, HLA-DP, -DQ and -DR. Human MHC class II antigens are transmembrane glycoproteins composed of an alpha chain (36 kDa) and a beta chain (27 kDa). They...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[HLA-Pan/2967R]
Molekulargewicht	33-35 kDa
UniProt	<a href="#">P04440</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody ELISA: For coating use Ab at 1-5 ug/mL order Ab without BSA Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes