

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **MUC2 (Mucin 2) (MLP/2970R), CF647 conjugate, 0.1mg/mL, Clone: [MLP/2970R], Rabbit, Monoclonal BOT-BNC472970-100**

Artikelname	MUC2 (Mucin 2) (MLP/2970R), CF647 conjugate, 0.1mg/mL, Clone: [MLP/2970R], Rabbit, Monoclonal
Artikelnummer	BOT-BNC472970-100
Hersteller Artikelnummer	BNC472970-100
Alternativnummer	BOT-BNC472970-100-100UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human MUC2 protein
Konjugation	CF647
Produktbeschreibung	This antibody recognizes a single glycoprotein of 520 kDa, identified as mucin 2 (MUC2). This MAb shows no cross-reaction with human milk fat globule membranes, MUC1, or MUC3. Its epitope has been defined as GTQTP (GlyThrGlnThrPro). Mucins are high m...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[MLP/2970R]

Molekulargewicht	520 kDa
UniProt	<a href="#">Q02817</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris-HCl buffer pH 10.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user