

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Neurofilament (NF-L) (Neuronal Marker) (NEFL/2983R), CF647 conjugate, 0.1mg/mL, Clone: [NEFL/2983R], Rabbit, Monoclonal BOT-BNC472983-100**

Artikelname	Neurofilament (NF-L) (Neuronal Marker) (NEFL/2983R), CF647 conjugate, 0.1mg/mL, Clone: [NEFL/2983R], Rabbit, Monoclonal
Artikelnummer	BOT-BNC472983-100
Hersteller Artikelnummer	BNC472983-100
Alternativnummer	BOT-BNC472983-100-100UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Bovine, Gallus, Human, Porcine, Rat
Immunogen	Recombinant full-length human NEFL protein
Konjugation	CF647
Produktbeschreibung	This MAb reacts with a 68 kDa protein, identified as light sub-unit of neurofilaments (NF-L). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerves, and sympathetic ganglion cells. Neur...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[NEFL/2983R]

Molekulargewicht	68 kDa
UniProt	<a href="#">P07196</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.25-0.5 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user