

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

HLA-DRB(LN-3 + HLA-DRB/1067), 0.2mg/mL, Clone: [LN-3 HLA-DRB/1067], Mouse, Monoclonal BOT-BNUB1208-100

Artikelname	HLA-DRB(LN-3 + HLA-DRB/1067), 0.2mg/mL, Clone: [LN-3 HLA-DRB/1067], Mouse, Monoclonal
Artikelnummer	BOT-BNUB1208-100
Hersteller Artikelnummer	BNUB1208-100
Alternativnummer	BOT-BNUB1208-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC, WB
Spezies Reaktivität	Human, Monkey
Immunogen	Activated human peripheral blood mononuclear cells (LN-3 & HLA-DRB/1067)
Produktbeschreibung	This MAb reacts with the beta-chain of HLA-DRB1 antigen, a member of MHC class II molecules. It does not cross react with HLA-DP and HLA-DQ. HLA-DR is a heterodimeric cell surface glycoprotein comprised of a 36 kDa alpha (heavy) chain and a 28 kDa be...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[LN-3 HLA-DRB/1067]

Molekulargewicht	28 kDa (beta chain)
UniProt	P01911
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user