

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin 8/18 (KRT8.18/1346), 0.2mg/mL, Clone: [KRT8.18/1346], Mouse, Monoclonal BOT-BNUB1346-100

Artikelname	Cytokeratin 8/18 (KRT8.18/1346), 0.2mg/mL, Clone: [KRT8.18/1346], Mouse, Monoclonal
Artikelnummer	BOT-BNUB1346-100
Hersteller Artikelnummer	BNUB1346-100
Alternativnummer	BOT-BNUB1346-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Cytoskeleton preparation from HeLa cells
Produktbeschreibung	Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). This MAb recognizes all simple epithelia including glandular epithelium, for example thyr...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[KRT8.18/1346]
Molekulargewicht	52.5 kDa (CK8), 45 kDa (CK18)

UniProt	P05787
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user