

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **MMP9 (rMMP9/1769), 0.2mg/mL, IgG1, Clone: [rMMP9/1769], Mouse, Monoclonal BOT-BNUB2176-100**

|                          |   |
|--------------------------|---|
| Artikelname              | MMP9 (rMMP9/1769), 0.2mg/mL, IgG1, Clone: [rMMP9/1769], Mouse, Monoclonal   |
| Artikelnummer            | BOT-BNUB2176-100  |
| Hersteller Artikelnummer | BNUB2176-100  |
| Alternativnummer         | BOT-BNUB2176-100-100UL  |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant full-length human MMP-9 protein   |
| Produktbeschreibung      | The matrix metalloproteinases (MMP) are a family of peptidase enzymes responsible for the degradation of extracellular matrix components, including collagen, gelatin, fibronectin, laminin and proteoglycan. Transcription of MMP genes is differentially... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.2 mg/mL   |
| Klon-Bezeichnung         | [rMMP9/1769]  |
| Molekulargewicht         | 92 kDa  |

|                        |   |
|------------------------|---|
| Isotyp                 | IgG1  |
| UniProt                | <a href="#">P14780</a>  |
| Puffer                 | PBS, 0.05% BSA, 0.05% azide   |
| Quelle                 | Animal  |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |