

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

MAP3K1 (Mitogen-Activated Protein Kinase Kinase Kinase 1)(2F6), 1mg/mL, Clone: [2F6], Mouse, Monoclonal BOT-BNUM0316-50

Artikelname	MAP3K1 (Mitogen-Activated Protein Kinase Kinase Kinase 1)(2F6), 1mg/mL, Clone: [2F6], Mouse, Monoclonal
Artikelnummer	BOT-BNUM0316-50
Hersteller Artikelnummer	BNUM0316-50
Alternativnummer	BOT-BNUM0316-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Partial recombinant MAP3K1 (aa1077-1176) (SKNSMTLDLNSSSKCDDSFSGCSSNSSNAVIPSDTVFTP- VEEKCRLDVNTELNSSIEDLLEASMPSSDTTVTFKSEVAVLSPEKAENDDTYK DDVNHNQK)
Produktbeschreibung	Mitogen-activated protein (MAP) kinase cascades are activated by various extracellular stimuli, including growth factors. The MEK kinases (also designated MAP kinase kinase kinases, MKKKs, MAP3Ks or MEKKs) phosphorylate and thereby activate the MEKs ...
Klonalität	Monoclonal
Konzentration	1 mg/mL

Klon-Bezeichnung	[2F6]
Molekulargewicht	195 kDa (intact), 80 kDa (cleaved)
UniProt	Q13233
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistochemistry (formalin-fixed): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris buffer with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Western Blot 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user