

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Cytokeratin 8(H1), 1mg/mL, IgG1, Mouse BOT-BNUM0334-50

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | Cytokeratin 8(H1), 1mg/mL, IgG1, Mouse                                                                                                                                                                                                                        |
| Artikelnummer            | BOT-BNUM0334-50                                                                                                                                                                                                                                               |
| Hersteller Artikelnummer | BNUM0334-50                                                                                                                                                                                                                                                   |
| Alternativnummer         | BOT-BNUM0334-50-50UL                                                                                                                                                                                                                                          |
| Hersteller               | Biotium                                                                                                                                                                                                                                                       |
| Wirt                     | Mouse                                                                                                                                                                                                                                                         |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | FC, IF, IHC, WB                                                                                                                                                                                                                                               |
| Spezies Reaktivität      | Human, Rat                                                                                                                                                                                                                                                    |
| Immunogen                | Cytoskeleton preparation containing cytokeratin 8                                                                                                                                                                                                             |
| Produktbeschreibung      | Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). CK8 is primarily found in the non-squamous epithelia and is present in majority of adenoc... |
| Konzentration            | 1 mg/mL                                                                                                                                                                                                                                                       |
| Molekulargewicht         | 52.5 kDa                                                                                                                                                                                                                                                      |
| Isotyp                   | IgG1                                                                                                                                                                                                                                                          |
| UniProt                  | <a href="#">P05787</a>                                                                                                                                                                                                                                        |
| Puffer                   | PBS, no BSA, no azide                                                                                                                                                                                                                                         |

| Quelle                 | Animal                                                                                                                                                                                                                                                                                                                                                         |
|------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Anwendungsbeschreibung | <p>Immunohistology formalin-fixed 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Immunofluorescence 1-2 ug/mL Optimal dilution for a specific application should be determined by user</p> |