

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### PCNA(PC10), 1mg/mL, Clone: [PC10], Mouse, Monoclonal BOT-BNUM0467-50

Artikelname	PCNA(PC10), 1mg/mL, Clone: [PC10], Mouse, Monoclonal
Artikelnummer	BOT-BNUM0467-50
Hersteller Artikelnummer	BNUM0467-50
Alternativnummer	BOT-BNUM0467-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IHC
Spezies Reaktivität	Drosophila, Gallus, Human, Monkey, Mouse, Porcine, Rat, Yeast, Zebrafish
Immunogen	Rat PCNA/Protein A fusion protein
Produktbeschreibung	Recognizes a non-histone protein of 36 kDa, which is identified as proliferating cell nuclear antigen (PCNA). It is also known as cyclin or polymerase delta auxiliary protein. Elevated expression of PCNA/cyclin has been shown in the nucleus during la...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[PC10]
Molekulargewicht	36 kDa

UniProt	<a href="#">P12004</a>
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user