

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD79a(JCB117), 1mg/mL, Clone: [JCB117], Mouse, Monoclonal BOT-BNUM0476-50

| | |
|--------------------------|---|
| Artikelname | CD79a(JCB117), 1mg/mL, Clone: [JCB117], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUM0476-50 |
| Hersteller Artikelnummer | BNUM0476-50 |
| Alternativnummer | BOT-BNUM0476-50-50UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | A synthetic peptide corresponding to aa 202-216 (GTQDVGSLNIADVQ) of human CD79a protein. |
| Produktbeschreibung | CD79 is a disulphide-linked heterodimer consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute... |
| Klonalität | Monoclonal |
| Konzentration | 1 mg/mL |
| Klon-Bezeichnung | [JCB117] |
| Molekulargewicht | 44 kDa |

| | |
|------------------------|---|
| UniProt | P11912 |
| Puffer | PBS, no BSA, no azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |