

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **ODC-1 (Ornithine Decarboxylase-1)(ODC1/486), 1mg/mL, Clone: [ODC1/486], Mouse, Monoclonal BOT-BNUM0486-50**

Artikelname	ODC-1 (Ornithine Decarboxylase-1)(ODC1/486), 1mg/mL, Clone: [ODC1/486], Mouse, Monoclonal
Artikelnummer	BOT-BNUM0486-50
Hersteller Artikelnummer	BNUM0486-50
Alternativnummer	BOT-BNUM0486-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human, Rat
Immunogen	Recombinant human ODC-1 protein
Produktbeschreibung	Recognizes a 53 kDa protein, identified as the Ornithine Decarboxylase (ODC-1). ODC is the initial and rate-limiting enzyme in the biosynthetic pathway of polyamines and is involved in the conversion of ornithine to putrescine. The biological activit...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[ODC1/486]
Molekulargewicht	53 kDa

UniProt	<a href="#">P11926</a>
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user