

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Myogenin(F5D), 1mg/mL, Clone: [F5D], Mouse, Monoclonal BOT-BNUM0632-50

| | |
|--------------------------|---|
| Artikelname | Myogenin(F5D), 1mg/mL, Clone: [F5D], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUM0632-50 |
| Hersteller Artikelnummer | BNUM0632-50 |
| Alternativnummer | BOT-BNUM0632-50-50UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Canine, Feline, Human, Mouse, Porcine, Rat |
| Immunogen | Rat myogenin peptide (aa 73-94) followed by rat myogenin recombinant fragment (aa30-224) (Epitope aa138-158) |
| Produktbeschreibung | Myogenin is a member of the MyoD family of myogenic basic helix-loop-helix (bHLH) transcription factors that also includes MyoD, Myf-5, and MRF4 (also known as herculinor Myf-6). MyoD family members are expressed exclusively in skeletal muscle and pl... |
| Klonalität | Monoclonal |
| Konzentration | 1 mg/mL |
| Klon-Bezeichnung | [F5D] |
| Molekulargewicht | 34 kDa |

| | |
|------------------------|---|
| UniProt | P15173 |
| Puffer | PBS, no BSA, no azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunoprecipitation: 0.5-1 ug/500 ug protein lysate Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1.0 ug/mL ELISA For coating, order Ab without BSA Optimal dilution for a specific application should be determined by user |