

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Smooth Muscle Actin(1A4), 1mg/mL, Clone: [1A4], Mouse, Monoclonal BOT-BNUM0665-50

| | |
|--------------------------|---|
| Artikelname | Smooth Muscle Actin(1A4), 1mg/mL, Clone: [1A4], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUM0665-50 |
| Hersteller Artikelnummer | BNUM0665-50 |
| Alternativnummer | BOT-BNUM0665-50-50UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC |
| Spezies Reaktivität | Bovine, Canine, Feline, Gallus, Goat, Guinea pig, Human, Monkey, Mouse, Porcine, Primate, Rabbit, Rat, Sheep |
| Immunogen | N-Terminal decapeptide of alpha smooth muscle isoform of actin and conjugated to KLH. |
| Produktbeschreibung | Actin is a major component of the cytoskeleton and is present in most cell types. This MAb is highly specific to actin from smooth muscles. Its epitope lies in the first four N-terminal amino acids. This MAb does not stain cardiac or skeletal muscle,... |
| Klonalität | Monoclonal |
| Konzentration | 1 mg/mL |
| Klon-Bezeichnung | [1A4] |

| | |
|------------------------|---|
| Molekulargewicht | 42 kDa |
| UniProt | P62736 |
| Puffer | PBS, no BSA, no azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |